# (8 points)

# 3. In each column, circle the appropriate characteristic for the data described.

(a) A geologist studies the <b>density</b> (grams per cubic cm) of different volcanic rocks.	Qualitative Quantitative	Discrete Continuous	Nominal Ordinal	Interval Ratio
(b) A department store records the <b>number</b> of customers that bought "pet rocks" in November	Qualitative Quantitative	Discrete Continuous	Nominal Ordinal	Interval Ratio
(c) An insurance company counts the <b>number</b> of accident claims that involve rocks hitting car windows.	Qualitative Quantitative	Discrete Continuous	Nominal Ordinal	Interval Ratio
(d) An agent prepares a list with the <b>names</b> of 37 rock bands in the Sacramento area.	Qualitative Quantitative	Discrete Continuous	Nominal Ordinal	Interval Ratio
(e) An earthquake investigator measures the <b>ages</b> of rocks (how many years since each rock was formed).	Qualitative Quantitative	Discrete Continuous	Nominal Ordinal	Interval Ratio
(f) Volcano watchers measure the <b>temperatures</b> of rocks in degrees Celsius to predict erruptions.	Qualitative Quantitative	Discrete Continuous	Nominal Ordinal	Interval Ratio

(8 points: 8 minutes)

1. For the data in Column A, circle the appropriate chacteristics in Column B and Column D. If the characteristic you circled in in Column B is "Quantitative," than circle the appropriate characteristic in Column C.

Column A	Column B	Column C	Colum	n D
Scientists measure the temperatures of Alaskan sled dogs in ºF.	Qualitative Quantitative	Continuous Discrete	Nominal Interval	Ordinal Ratio
The names of Alaskan sled dogs that live longer than 7 years.	Qualitative Quantitative	Continuous Discrete	Nominal Interval	Ordinal Ratio
The number of times that "Fred," an Alaskan sled dog, turns all the way around before lying down to sleep.	Qualitative Quantitative	Continuous Discrete	Nominal Interval	Ordinal Ratio
The lengths of the hairs on the tails of Alaskan sled dogs.	Qualitative Quantitative	Continuous Discrete	Nominal Interval	Ordinal Ratio

(7 points: 10 minutes)

3. For each situation below, select the appropriate statistical term from the list provided and write it in the blank next to the description or situation. Choose the term that is best connected to the underlined text in the description or situation.

Terms:	1. randomization	5. placebo	
	2. replication	6. block	
	3. confounding	7. experimental unit	
	4. blinding	8. treatment	
A.		Each person included in the study be	alonged to a group of other
<u> </u>		<u>people</u> in the study who had similar p	
		similar lifestyles.	
B.		A scientist prepares experimental ma	aterial in groups of similar
-		units. Within each group, each expen	
		to a treatment so that all possible pai	irings of treatments and
		units are equally likely to occur.	
С.		A flower-growing company tests 100	rose bushes, each in its own
		pot. Each pot (or each bush) is treate	ed with one of two insect
		poisons, where the poison is chosen	at random.
D.		A veterinarian (animal doctor) works	
		compare the effectiveness of vitamin	••
		diet. Ten horse owners feed the hors	
		with supplement B. And, ten use sup	oplement C, which neither
		harms nor helps horses.	
E		An experiment was done to see how	
		to ozone, a pollutant that irritates lun	
		children had more asthma symptoms fewer symptoms when ozone was low	-
		on Sunday and low on Monday, so a	· · · · · · · · · · · · · · · · · · ·
		children's response might be due to	
		due to ozone.	<u> </u>
F.		A political consultant tests three diffe	erent TV adds for possible
_		use in the campaign. One add is test	ed in Sab Francisco, another
		in Fresno, and another in Orange Co	unty. An important feature
		is lacking in this experiment.	
G		Ten cars (with their drivers) will be us	-
		for mileage (miles per gallon). Techn	
		so <u>the driver will not know gasoline "</u>	<u>'A" or "B" is being used</u> .
н		Ten cars (with their drivers) will be us	
		for mileage (miles per gallon). Techn	
		so the driver will not know gasoline "	<u>'A" or "B"</u> is being used.

(4 points: 5 minutes)

2. A real estate developer is interested in the values of homes in California. In particular, he/she wants to know whether the median value of all homes in California is now more than \$200,000. To explore this question, 500 in California homes are selected at random and their current values are determined. The median value of the 500 homes is \$212,800, which is more than \$200,000.

Use the information in the "story" to answer the following:

(a) What is the population of interest?

(c) What statistic was used?

(d) What parameter was the parameter of interest?

(d) Was a census or a sample used in the work?

(e) How do you know whether a census or a sample was used?

(6 points, 12 minutes)

- 5. Identify each situation below as RANDOM, STRATIFIED, SYSTEMMATIC, CLUSTER, or CONVENIENCE sampling.
  - (a) A chain of pizza restaurants wants to know the preferences of their customers for different kinds of pizza. The chain arranges for a study to select 200 households at random. At each household, the preferences of all residents are determined.
  - (b) A chain of pizza restaurants (stores) wants to know whether customer preferences are the same in urban, suburban, and rural areas. They select 10 urban stores at random, 8 suburban stores at random, and 6 rural stores at random. The sales of each type of pizza at the selected stores are compared.
  - (c) The owner of a music store believes his customers eat lots of pizza. Therefore, he goes to local pizza restaurants and asks the restaurant's customers what kinds of CD's they like to buy. Some of the people are glad to tell him their music preferences but others tell him to go away.
  - (d) A factory makes frozen pizzas for grocery stores. The quality control unit at the factory tests every 500th pizza to ensure that the product meets the required standards for quality.
  - (e) The federal covernment wants to know how long workers at pizza restaurants keep their jobs. By using data from income tax records, the government prepares a sample of 2000 pizza workers. Every pizza worker has an equal chance of being in the sample. The length of time each worker worked at their job is discovered.

# 1. Identify each of the following "sampling" situations as RANDOM, STRATIFIED, SYSTEMMATIC, CLUSTER, or CONVENIENCE sampling.

A marketing company conducts a survey of product preferences. To make sure that their results are useful to many customers, they select random samples of 500 individuals from each of 10 age groups.

A quality control manager needs to report on product quality, so he directs the testing technicians to get 100 items out of the stack of boxes on the shipping platform and test them.

A supervisor monitors the accuracy of every 50<sup>th</sup> telephone call handled by customer service agents.

A chain of pizza restaurants decides to examine customer preferences. A list of tables at all of the chain's restaurants is prepared, and tables are selected at random. All customers that sit at each selected table are asked whether they like the pizza theyare eating.

The Highway Patrol wants to test the value of their new "traffic school." They select 400 people convicted of drunk driving in the last two years in a way that gives all such convicts an equal chance of being selected. Half of those selected are assigned to the old "traffic school" and half to the new "traffic school."

(e)

(b)

(c)

(a)

(d)

# (6 points, 6 minutes)

1. Use the data below to determine the frequencies, relative frequencies, cumulative frequencies, and cumulative relative frequencies for the frequency table.

	ľ							Relative
Data		Class	Limits			Relative	Cumulative	Cumulative
		Upper	Lower	Tally	Frequency	Frequency	Frequency	Frequency
3587.4								
2335.4		1000	2000					
1735.0								
6091.5		3000	4000					
4184.3								
3423.2		5000	6000					
2833.5								
3522.3		7000	8000					
5052.0								

5052.0 7310.6 5885.3

3311.1

(8 points)

2. Complete the columns in the "Frequency Distribution" table using the data values given below, and determine the class width.

			equency Dis				
Class Lower	Limits Upper	Tally	Frequency	Cumulative Frequency		Class Midpoint	Clas Bound
1.4	1.6						
1.7	1.9						
2.0	2.2						
2.3	2.5					-	

**Frequency Distribution** 

Г	Deter	2.40	0.00	0.40	4.04	2.4.4	0.04
	Data:	2.16	2.33	2.43	1.84	2.14	2.24
		1.98	2.05	1.64	2.19	2.08	2.22
		1.56					

Clas Midpo	_	Class Boundary
		1

Class width =

3. For the data in this problem, calculate the following sample statistics: RANGE, STANDARD DEVIATION ,and VARIANCE. You may use your calculators or do the calculations "by hand".

Data: 11 8 16 20 6 8	Range =
	Variance =

(2 points)

E


\_

4. Without calculating it directly, estimate the standard deviation for the data in problem 3 above. (Show how you did the estimation)

Estimate of s = \_\_\_\_\_

### (14 points, 10 minutes)

3. Use the data below to determine the value of each statistic in parts (a) through (g). Write an expression for each statistic or describe how it is calculated (do NOT describe how to use the calculator).



(4 minutes)

- 7. Plant species "A" has leaves that average 8 centimeters long with a standard deviation of 0.8 centimeters. Plant species "B" has flowers with an average width of 2 centimeters with a standard deviation of 0.25 centimeters. Which of the two plants below is the most unusual? Circle (a) or (b) and show why!
  - (a) A plant from species "A" has leaves that are 6 centimeters long.
  - (b) A plant from species "B" has flowers that are 2.5 centimeters wide.

(4 minutes)

8. A plant species has leaves that average 8 centimeters long with a standard deviation of 0.8 centimeters. The distribution of leaf lengths is approximately "bell-shaped". What percentage of the leaves from plants of this species should be between 5.6 cm and 10.6 cm long? Show why! (6 points)5. Provide the "z-scores" for each situation.

	μ	σ	x	z
(a)	60	20	88	
(b)	60	5	52	
(c)	60	4	68	

(d) If the distribution is "Bell-shaped", which "X" value is the most unusual?

(e) Approximately what percent of the data are closer to the mean than the X value in your answer for part (d)? [Be sure to explain how you get your estimate]

(3 points)6. For the data below, what is the percentile (P?) of the value 486 ?

736	486	26	360	186	758	548

(3 points)

 7. For the data below (given in sorted order), what is the 27<sup>th</sup> percentile? [note: There are 60 values in the dataset, five values per row]

1767	1772	1777	1784
1810	1837	1861	1896
1945	1967	1973	1985
2007	2008	2039	2060
2098	2126	2147	2172
2208	2210	2223	2248
2272	2311	2316	2330
2354	2380	2403	2423
2475	2481	2492	2518
2585	2596	2630	2656
2706	2708	2712	2729
2775	2791	2805	2821
	1810 1945 2007 2098 2208 2272 2354 2475 2585 2706	1810 1837   1945 1967   2007 2008   2098 2126   2208 2210   2272 2311   2354 2380   2475 2481   2585 2596   2706 2708	181018371861194519671973200720082039209821262147220822102223227223112316235423802403247524812492258525962630270627082712

- 4. The data at the bottom of the page are in seven rows of ten values and the values are sorted from smallest to largest. Use these data to answer parts (a) and (b) of this question.
  - (4 points, 4 minutes)
  - (a) What value represents to 83rd percentile, P83?

(4 points, 4 minutes)

(b) What percentile, P?, is represented by the value 222?

200	210	212	220	222	222	226	228	234	240
244	252	255	259	264	271	276	286	292	299
299	305	310	314	317	327	336	340	343	347
355	360	365	366	367	373	381	387	387	392
402	402	405	410	413	420	420	426	436	440
441	447	454	463	466	468	474	477	481	485
492	494	502	508	516	522	523	524	524	524

(6 points)

9. Use the information in the table to answer the probability questions (a) - (d).

Favorite				
Basketball Team	Arizona California		Oregon	Total
Phoenix Suns	160	60	30	250
Sacramento Kings	20	270	10	300
Portland Trailblazers	20	70	160	250
Total	200	400	200	800

(a) What is the probability that a randomly selected person from this survey would be from California?

(b) What is the probability that a randomly selected person's favorite team would be the Sacramento Kings given that the person is from Arizona?

(c) What is the probability that a randomly selected person would be from Oregon or have the Portland Trailblazers as their favorite team?

(d) What is the probability that a randomly selected person's favorite team would be the Sacramento Kings given that the person is from Oregon?

(e) Use your answers from part (b) and part (d) to demonstrate whether or not a person's preference of team is independent of their home state [1 point extra credit - do this last].

(5 points)

- 10. A collection of 20 dice includes 4 bad dice (not balanced correctly) and 16 dice that are good (balanced correctly).
  - (a) What is the probability that a random sample of four of the 20 dice (without replacement) will have the exact sequence : "Good", "Good", "Good", "Bad" ?

(b) What is the probability that a random sample of four of the 20 dice (without replacement) will have at least one "Bad" die?

(2 points)

11. At an athletic contest, performers are judged by three judges. Each judge scores each performer as a 1, 2, 3, 4, 5, or a "perfect" 6. How many different sequences of scores from Judge1, Judge2, and Judge 3 are possible?

Example:	2	2	5	
-	Judge 1	Judge 2	Judge 3	

(2 points)

12. In an office, there are six workers and all of them have equal seniority. If there are six desks, how many ways can the six workers be assigned to the six desks?

(2 points)

13. In the same office as problem 12, the manager decides to assign the six workers to the six desks randomly. What is the probability that the random assignments result in the workers being assigned to the desk in the same order as their ages from oldest to youngest?

# (3 points)

14. A rich man has 14 close friends. He decides to invite three of these friends to join him on a voyage around the world. If the man selects his three traveling companions at random, how many different traveling groups could be formed?

6. Use the data in the following table to answer parts (a) through (e).

(2 points, 1 minutes)		Employment Type for People in Sample					
(a) What is the probability that	State	Farm	Factory	Gov'ment	Service	Total	
a randomly selected person from this sample will be a factory worker?	Arizona	21	36	14	29	100	
	California	28	32	11	49	120	
	Nevada	19	41	13	7	80	
	Oregon	24	37	14	15	90	
	Washington	15	36	13	36	100	
	Total	107	182	65	136	490	

(2 points, 2 minutes)

(b) What is the probability that a person randomly selected form this sample will be from California given that the person works in the Service industry?

(3 points, 3 minutes)

(c) What is the probability that a person randomly selected form this sample will be from Oregon or work in government?

(2 points, 2 minutes)

(d) What is the probability that a person randomly selected form this sample will be from Arizona or Washington?

### (3 points: 4 minutes)

3. The local Police force has been accused of being insensitive to racial differences among the officers in the force. To answer the charges and to try to increase the multi-cultural awareness of officers, all the teams of two partners are mixed up and reassigned as new teams. If there are 16 officers on the force, how many different teams of two officers could possibly be formed?

(3 points: 4 minutes)

4. On the same police force as used in Problem 3, the 16 officers include 8 White officers, 4 Latino officers, 2 Black officers, and 2 Asian officers. If the entire department forms one line to take a picture, how many different racial sequences would be possible from left to right?

(3 points: 4 minutes)

5. A police officer gets stuck one day with an unwelcome job – defusing a bomb – because the bomb squad's truck will not start. The officer sees a red wire, a white wire, a green wire, and a black wire. To stop the bomb, two wires must be cut. Because the officer has no better idea what to do, he decides to cut one wire and then another at random. What is the sample space for the officer's experiment? Also, what is the probability that the officer randomly selects the correct choice of two wires to cut to stop the bomb from exploding?

(6 points: 6 minutes)

- 11. The "Bomb Squad" must disable a terrorist bomb. This bomb has 6 wires. Three wires must be cut, but only the right 3 wires. All the wires are RED. For this situation, answer part (a) and part (b).
  - (a) How many sets of three wires are possible for the Bomb Squad to choose?
  - (b) If the Bomb Squad must choose the right 3 in the right order "OR ELSE!", and they must choose at random, what is the probability the bomb does not explode?

(2 points)

11. At an athletic contest, performers are judged by three judges. Each judge scores each performer as a 1, 2, 3, 4, 5, or a "perfect" 6. How many different sequences of scores from Judge1, Judge2, and Judge 3 are possible?

(2 points)

12. In an office, there are six workers and all of them have equal seniority. If there are six desks, how many ways can the six workers be assigned to the six desks?

(5 points)

10. A box of 80 spark plugs for car engines includes 70 good ones and 10 bad ones.

(a) If two spark plugs are selected at random (without replacement) what is the probability that the first will be bad and the second will also be bad?

(b) What is the probability that a random sample of six spark plugs (without replacement) will have at least one "Bad" one?

(2 points)

11. Olympic ice skaters are judged by three judges. If each judge can score each performer as a 1, 2, 3, 4, 5, or a "perfect" 6 (no decimals), how many different sequences of scores from the six judges are possible?

Example:	2	2 5		2	2	5	
	Judge 1	Judge 2	Judge 3	Judge 1	Judge 2	Judge 3	

(2 points)

12. A carnival has 13 different "game" booths. The manager has to assign places along the "midway". How many different ways can the manager arrange the booths along one side of the path that customers walk?

(2 points)

13. For the carnival in problem 12 (above), the manager has a problem: there is only room for 9 of the 13 game booths at the next town. How many ways could the manager choose the 9 booths that will set up and operate when the show arrives at the next town?

#### (4 points)

16. A business analyst determines that a new business will earn \$100,000 in profits if it is successful, but it will lose \$20,000 if it fails. If the probability of success is 0.85 and the probability of failure is 0.15, what is the expected value of a decision to start the new business?

(4 points)

17. A manufacturing process produces silicon chips. The frequency (at random) of defective chips is 0.3. In a random sample of 6 silicon chips, what is the probability that the number of defective chips will be zero or one? [Assume the sample comes from a population of millions of chips so the selections are independent and have the same probability of being defective.]

For full credit, calculate the probability. For almost full credit, set up the full solution to the problem. (6 points)

15. For each discrete probability distribution below, calculate the mean, variance, and standard deviation. [Use extra columns however you want to.]

(a)	х	P(x)			
	0	0.2			
	1	0.7			
	2	0.1			

(b)	x	P(x)			
	10	0.2			
	20	0.2			
	30	0.4			

#### (4 points)

16. A business analyst determines that a new business will earn \$100,000 in profits if it is successful, but it will lose \$20,000 if it fails. If the probability of success is 0.85 and the probability of failure is 0.15, what is the expected value of a decision to start the new business?

(4 points)

17. A manufacturing process produces silicon chips. The frequency (at random) of defective chips is 0.3. In a random sample of 6 silicon chips, what is the probability that the number of defective chips will be zero or one? [Assume the sample comes from a population of millions of chips so the selections are independent and have the same probability of being defective.]

For full credit, calculate the probability. For almost full credit, set up the full solution to the problem.

## (5 minutes)

11. The number of boys in 168 births is distributed binomially with n = 168 and p = 0.50. If there are 70 boys in a randomly selected set of 168 births, is that result unusual?

(8 minutes)

12. An investment opportunity has two possible outcomes. The investor may earn \$100 or the investor may lose \$20. If the probability of earning \$100 is 0.26 and the probability of losing \$20 is 0.74, what is the expected value of the investment.